

The coverage probability of confidence intervals in regression after a preliminary F test

PAUL KABAILA^{1*} AND DAVIDE FARCHIONE²

1. Department of Mathematics and Statistics, La Trobe University, Victoria 3086, Australia

2. School of Mathematics and Statistics, University of South Australia, South Australia 5095, Australia

* Author to whom correspondence should be addressed.

E-mail: P.Kabaila@latrobe.edu.au, Facsimile: 61 3 9479 2466, Telephone: 61 3 9479 2594

Abstract

Consider a linear regression model with regression parameter $\beta = (\beta_1, \dots, \beta_p)$ and independent normal errors. Suppose the parameter of interest is $\theta = a^T \beta$, where a is specified. Define the s -dimensional parameter vector $\tau = C^T \beta - t$, where C and t are specified. Suppose that we carry out a preliminary F test of the null hypothesis $H_0 : \tau = 0$ against the alternative hypothesis $H_1 : \tau \neq 0$. It is common statistical practice to then construct a confidence interval for θ with nominal coverage $1 - \alpha$, using the same data, based on the assumption that the selected model had been given to us *a priori* (as the true model). We call this the naive $1 - \alpha$ confidence interval for θ . This assumption is false and it may lead to this confidence interval having minimum coverage probability far below $1 - \alpha$, making it completely inadequate. Our aim is to compute this minimum coverage probability. It is straightforward to find an expression for the coverage probability of this confidence interval that is a multiple integral of dimension $s + 1$. However, we derive a new elegant and computationally-convenient formula for this coverage probability. For $s = 2$ this formula is a sum of a triple and a double integral and for all $s > 2$ this formula is a sum of a quadruple and a double integral. This makes it easy to compute the minimum coverage probability of the naive confidence interval, irrespective of how large s is. A very important practical application of this formula is to the *analysis of covariance*. In this context, τ can be defined so that H_0 expresses the hypothesis of “parallelism”. Applied statisticians commonly recommend carrying out a preliminary F test of this hypothesis. We illustrate the application of our formula with a real-life analysis of covariance data set and a preliminary F test for “parallelism”. We show that the naive 0.95 confidence interval has minimum coverage probability 0.0846, showing that it is completely inadequate.

Keywords: analysis of covariance, naive confidence interval, preliminary F test, test for parallelism.

1. Introduction

Consider the linear regression model $Y = X\beta + \varepsilon$, where Y is a random n -vector of responses, X is a known $n \times p$ matrix with linearly independent columns, β is an unknown parameter p -vector and $\varepsilon \sim N(0, \sigma^2 I_n)$ where σ^2 is an unknown positive parameter. Suppose that the parameter of interest is $\theta = a^T \beta$ where a is a given p -vector ($a \neq 0$). We seek a $1 - \alpha$ confidence interval for θ .

Let the s -dimensional parameter vector τ be defined to be $C^T \beta - t$ where C is a specified $p \times s$ matrix ($s < p$) with linearly independent columns and t is a specified s -vector. Suppose that a does not belong to the linear subspace spanned by the columns of C . Also suppose that we carry out a preliminary F test of the null hypothesis $H_0 : \tau = 0$ against the alternative hypothesis $H_1 : \tau \neq 0$. It is then common statistical practice to construct a confidence interval for θ with nominal coverage $1 - \alpha$, using the same data, based on the assumption that the selected model had been given to us *a priori* (as the true model). We call this the naive $1 - \alpha$ confidence interval for θ . In Section 2, we provide a convenient description of this confidence interval. This assumption is false and it can lead to the naive $1 - \alpha$ confidence interval having minimum coverage probability far below $1 - \alpha$, making it completely inadequate. Our aim is to compute this minimum coverage probability. For $s = 1$, the preliminary F test is equivalent to a t test. The case of a single preliminary t test has been dealt with by Kabaila and Giri (2009, Theorem 3). So, in the present paper, we restrict attention to the case that $s > 1$.

Straightforward application of the methodology of Farchione (PhD thesis, 2009, Section 5.7) leads to an expression for the coverage probability of the naive $1 - \alpha$ confidence interval, for a given value of an s -dimensional parameter vector, that is a multiple integral of dimension $s + 1$. Finding the minimum coverage probability using this formula becomes increasingly cumbersome as s increases due to both the need to (a) evaluate multiple integrals of dimension $s + 1$ and (b) the need to search for the minimum over a space of dimension s .

In Section 3, by a careful consideration of the geometry of the situation, we derive a new elegant and computationally-convenient formula for the coverage probability of this confidence interval for given parameter values. For $s = 2$ this formula is a sum of a triple and a double integral and for all $s > 2$ this formula is a sum of a quadruple and a double integral. This formula also shows that the coverage probability is a function of a two-dimensional parameter vector, irrespective of how

large s is. This makes it easy to compute the minimum coverage probability of the naive confidence interval, irrespective of how large s is. Another important aspect of this formula is that it can be used to delineate general categories of a , C and X for which the naive confidence interval has poor coverage properties.

A very important practical application of this formula is to the *analysis of covariance*. In this context, τ can be defined so that H_0 expresses the null hypothesis of “parallelism”. In the applied statistics literature on the analysis of covariance it is commonly recommended that a preliminary F test of the null hypothesis of “parallelism” be carried out. See, for example, Kuehl (2000, p.563), Milliken and Johnson (2002, pp. 14 – 17) and Freund et al (2006, pp. 363 – 368). For an analysis of covariance, we can choose a so that the parameter θ is the difference in expected responses for two specified treatments, for the same specified values of the covariates.

In Section 4, we illustrate the application of the results of the paper with a real-life *analysis of covariance* data set and a preliminary F test for “parallelism”. We define θ to be (expected response to treatment 1) – (expected response to treatment 2), evaluated at the same specified value of the covariate. We show that the naive 0.95 confidence interval for θ has minimum coverage probability 0.0846, for this specified value of the covariate. This shows that this confidence interval is completely inadequate, for this specified value of the covariate.

2. Description of the naive confidence interval

In this section we provide a convenient description of the naive $1 - \alpha$ confidence interval constructed after the preliminary F test. Let $\hat{\beta}$ denote the least squares estimator of β . Define $R(\beta) = (Y - X\beta)^T(Y - X\beta)$. Let $m = n - p$. Define $\hat{\Sigma}^2 = R(\hat{\beta})/m = (Y - X\hat{\beta})^T(Y - X\hat{\beta})/m$. Also, define $\hat{\Theta} = a^T\hat{\beta}$ and $\hat{\tau} = C^T\hat{\beta} - t$. We suppose that the columns of the matrix C are linearly independent. We also suppose that a does not belong to the linear subspace spanned by the columns of C . Now define the $(s + 1) \times (s + 1)$ matrix

$$V = \begin{bmatrix} v_{11} & v_{21}^T \\ v_{21} & V_{22} \end{bmatrix} = \frac{1}{\sigma^2} E \left(\begin{bmatrix} \hat{\Theta} - \theta \\ \hat{\tau} - \tau \end{bmatrix} \begin{bmatrix} \hat{\Theta} - \theta & (\hat{\tau} - \tau)^T \end{bmatrix} \right).$$

Note that $v_{11} = a^T(X^TX)^{-1}a$, $v_{21} = C^T(X^TX)^{-1}a$ and $V_{22} = C^T(X^TX)^{-1}C$.

Define β^* to be the value of β minimizing $R(\beta)$ subject to the restriction that

$\tau = C^T \beta - t = 0$. As is well known (see e.g. Graybill, 1976, p.222)

$$\begin{aligned}\beta^* &= \hat{\beta} - (X^T X)^{-1} C (C^T (X^T X)^{-1} C)^{-1} \hat{\tau} \\ R(\beta^*) &= R(\hat{\beta}) + (\hat{\beta} - \beta^*)^T (X^T X) (\hat{\beta} - \beta^*).\end{aligned}$$

The standard test statistic for testing $H_0 : \tau = 0$ against $H_1 : \tau \neq 0$ is

$$F = \frac{(\hat{\beta} - \beta^*)^T X^T X (\hat{\beta} - \beta^*) / s}{\hat{\Sigma}^2} = \frac{\hat{\tau}^T V_{22}^{-1} \hat{\tau} / s}{\hat{\Sigma}^2}.$$

This test statistic has an $F_{s,m}$ distribution under H_0 . Suppose that we reject H_0 when $F > \ell$ and accept H_0 otherwise, where ℓ is a specified positive number.

Define $\Theta^* = a^T \beta^*$. Also define the quantile $t(m)$ by the requirement that $P(-t(m) \leq T \leq t(m)) = 1 - \alpha$ for $T \sim t_m$. The naive $1 - \alpha$ confidence interval for θ is obtained as follows.

Suppose that $F > \ell$. The confidence interval is constructed on the assumption that $\tau = 0$ is not necessarily true. In this case, the naive $1 - \alpha$ confidence interval is the usual $1 - \alpha$ confidence interval for θ based on fitting the full model,

$$I = [\hat{\Theta} - t(m) \sqrt{v_{11}} \hat{\Sigma}, \hat{\Theta} + t(m) \sqrt{v_{11}} \hat{\Sigma}].$$

Now suppose that $F \leq \ell$. The confidence interval is constructed on the assumption that $\tau = 0$. If $\tau = 0$ then $R(\beta^*)/\sigma^2 \sim \chi_{m+s}^2$ and $\text{Var}(\Theta^*) = \sigma^2(v_{11} - v_{21}^T V_{22}^{-1} v_{21})$. Note that Θ^* and $R(\beta^*)$ are independent random variables. We use the notation $[a \pm b]$ for the interval $[a - b, a + b]$ ($b > 0$). In this case, the naive $1 - \alpha$ confidence interval for θ is

$$\begin{aligned}J &= \left[\Theta^* \pm t(m+s) \sqrt{\frac{R(\beta^*)}{m+s}} \sqrt{v_{11} - v_{21}^T V_{22}^{-1} v_{21}} \right] \\ &= \left[\Theta^* \pm t(m+s) \sqrt{\frac{R(\hat{\beta}) + \hat{\tau}^T V_{22}^{-1} \hat{\tau}}{m+s}} \sqrt{v_{11} - v_{21}^T V_{22}^{-1} v_{21}} \right].\end{aligned}\quad (1)$$

3. The coverage probability of the naive confidence interval

Define $b = v_{11}^{-1/2} V_{22}^{-1/2} v_{21}$ and $W = \hat{\Sigma}/\sigma$. Let f_W denote the probability density function of W . Define $\|b\| = \sqrt{b_1^2 + \dots + b_s^2}$. Thus

$$\begin{aligned}\|b\|^2 &= v_{11}^{-1} v_{21}^T V_{22}^{-1} v_{21} \\ &= \frac{a^T (X^T X)^{-1} C (C^T (X^T X)^{-1} C)^{-1} C^T (X^T X)^{-1} a}{a^T (X^T X)^{-1} a}.\end{aligned}$$

Since $\text{Var}(\Theta^*) = \sigma^2(v_{11} - v_{21}^T V_{22}^{-1} v_{21}) \geq 0$, $\|b\| \in [0, 1]$. The assumption that the vector a does not belong to the linear subspace spanned by the columns of C implies that $\|b\| > 0$. So, we may assume that $\|b\| \in (0, 1]$. Now define

$$i(x, w; \|b\|) = P(-t(m)w + x \leq Z \leq t(m)w + x),$$

where $Z \sim N(0, 1 - \|b\|^2)$, and

$$j(x, y, w; \|b\|) = P\left(x - t(m+s)\sqrt{\frac{mw^2 + y}{m+s}}\sqrt{1 - \|b\|^2} \leq Z \leq x + t(m+s)\sqrt{\frac{mw^2 + y}{m+s}}\sqrt{1 - \|b\|^2}\right).$$

Define f_R to be the probability density function of $\sqrt{R^2}$ when $R^2 \sim \chi_s^2$. Let $B(a, b)$ denote the beta function. Define the probability density function f_{T_1} to be

$$f_{T_1}(t_1) = \begin{cases} \frac{\pi}{B(\frac{1}{2}, \frac{s-1}{2})} \sin^{s-2}(\pi t_1) & \text{for } 0 \leq t_1 \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

For $s \geq 3$, define the probability density function f_{T_2} to be

$$f_{T_2}(t_2) = \begin{cases} \frac{\pi}{B(\frac{1}{2}, \frac{s-2}{2})} \sin^{s-3}(\pi t_2) & \text{for } 0 \leq t_2 \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Let $\gamma = (1/\sigma)V_{22}^{-1/2}\tau$. Define f_Q to be the probability density function of a noncentral chi squared distribution with s degrees of freedom and noncentrality parameter $\|\gamma\|^2$. Also define

$$d(t_1, r; s, \|\gamma\|) = \begin{cases} \|\gamma\|^2 + 2\|\gamma\|r \cos(2\pi t_1) + r^2 & \text{for } s = 2 \\ \|\gamma\|^2 + 2\|\gamma\|r \cos(\pi t_1) + r^2 & \text{for } s \geq 3. \end{cases}$$

Define the unit vector $u_b = (1/\|b\|)b$. When $\|\gamma\| > 0$, define $u_\gamma = (1/\|\gamma\|)\gamma$ and then $\psi = u_b^T u_\gamma$. Also define $\psi = 1$ when $\|\gamma\| = 0$. Now, when $\|\gamma\| > 0$, define

$$k(t_1; \psi) = \psi \cos(2\pi t_1) + \sqrt{1 - \psi^2} \sin(2\pi t_1)$$

$$k(t_1, t_2; s, \psi) = \begin{cases} \psi \cos(\pi t_1) + \sqrt{1 - \psi^2} \sin(\pi t_1) \cos(2\pi t_2) & \text{for } s = 3 \\ \psi \cos(\pi t_1) + \sqrt{1 - \psi^2} \sin(\pi t_1) \cos(\pi t_2) & \text{for } s \geq 4. \end{cases}$$

The following is the main result of the paper.

Theorem 1. *The coverage probability of the naive $1 - \alpha$ confidence interval for θ is $P(\theta \in I, F > \ell) + P(\theta \in J, F \leq \ell)$. A computationally-convenient expression for the second term in this sum is*

$$P(\theta \in J, F \leq \ell) = \int_0^\infty \int_0^{s\ell w^2} j(\|b\| \|\gamma\| \psi, q, w; \|b\|) f_Q(q) f_W(w) dq dw \quad (2)$$

and computationally-convenient expressions for $P(\theta \in I, F > \ell)$ are as follows. Let $u = \sqrt{d(t_1, r; s, \|\gamma\|)/s\ell}$. For $s = 2$,

$$P(\theta \in I, F > \ell) = \int_0^1 \int_0^\infty \int_0^u i(-\|b\| r k(t_1; \psi), w; \|b\|) f_W(w) f_R(r) dw dr dt_1 \quad (3)$$

For $s \geq 3$ and $\|\gamma\| > 0$, $P(\theta \in I, F > \ell)$ is equal to

$$\int_0^1 \int_0^1 \int_0^\infty \int_0^u i(-\|b\| r k(t_1, t_2; s, \psi), w; \|b\|) f_W(w) f_R(r) f_{T_1}(t_1) f_{T_2}(t_2) dw dr dt_1 dt_2 \quad (4)$$

For $s \geq 3$, $\|\gamma\| > 0$ and $\psi \in \{-1, 1\}$,

$$P(\theta \in I, F > \ell) = \int_0^1 \int_0^\infty \int_0^u i(-\|b\| r \cos(\pi t_1), w; \|b\|) f_W(w) f_R(r) f_{T_1}(t_1) dw dr dt_1$$

For $s \geq 3$ and $\|\gamma\| = 0$,

$$P(\theta \in I, F > \ell) = \int_0^1 \int_0^\infty \int_0^{r/\sqrt{s\ell}} i(-\|b\| r \cos(2\pi t_1), w; \|b\|) f_W(w) f_R(r) dw dr dt_1$$

Note that for given $\|b\|$ (which is determined by a , C and X) and m , s , ℓ and α , the coverage probability of the naive $1 - \alpha$ confidence interval is a function of $(\|\gamma\|, \psi)$.

The proof of this theorem is presented in Appendix A.

The formulas given in Theorem 1 have three attractive features. The first of these is that, irrespective of how large s is, these formulas involve, at most, a 4-dimensional integral. The second of these features is that the numerical evaluation of these integrals, reviewed in Appendix B, is very straightforward. Thirdly, for given m , s , ℓ , α and $\|b\|$, the coverage probability of the naive confidence interval is a function of the two-dimensional parameter vector $(\|\gamma\|, \psi)$. These features make it is easy to compute the minimum coverage probability of the naive $1 - \alpha$ confidence interval for given m , s , ℓ , α and $\|b\|$. Finally, Theorem 1 has the following corollary

Corollary 1. *For given m , s , ℓ and α , the infimum over $(\|\gamma\|, \psi)$ of the coverage probability of the naive $1 - \alpha$ confidence interval for θ is a function of $\|b\|$.*

For the numerical example described in the next section, $m = 4$, $s = 4$, $\ell = 6.5914$ (corresponding to a 0.05 level of significance of the preliminary F test) and $\alpha = 0.05$. For these values of m , s and ℓ , the minimum coverage probability of the naive 0.95 confidence interval, as a function of $\|b\| \in (0, 1]$, is as shown in Figure 1. All of the computations presented in the present paper were performed with programs written in MATLAB, using the optimization and statistics toolboxes. We note the decrease in the minimum coverage probability of this naive confidence interval with increasing $\|b\|$. We can use Corollary 1 to delineate general categories of a , C and X (via their relationship to $\|b\|$) for which this naive 0.95 confidence interval has poor coverage properties. Specifically, this naive confidence interval will have poor coverage properties for those values of a , C and X such that

$$\|b\| = \sqrt{\frac{a^T (X^T X)^{-1} C (C^T (X^T X)^{-1} C)^{-1} C^T (X^T X)^{-1} a}{a^T (X^T X)^{-1} a}}$$

is greater than 0.7, say.




Figure 1: Plot of the minimum coverage probability of the naive 0.95 confidence interval, as a function of $\|b\| \in (0, 1]$, for $m = 4$, $s = 4$, $\ell = 6.5914$ and $\alpha = 0.05$.

4. Application to a real-life data set

In this section we consider the real-life *analysis of covariance* data set due to Chin et al (1994) and analysed by Yandell (1997, Chapter 17), who makes this data available at the website <http://www.stat.wisc.edu/~yandell/pda/>. This data is listed in the Table 1. It consists of the observed response (weight gain) for a given treatment and value of the covariate (feed intake). There are 4 possible treatments, numbered 1 to 4.

treatment	weight gain	feed intake
1	1416.1	2451.75
1	1447.0	2546.00
1	1509.6	2657.00
2	1497.8	2452.10
2	1469.9	2404.90
2	1469.4	2479.90
3	1510.1	2788.70
3	1423.0	2655.50
3	1295.9	2366.40
4	1354.8	2578.80
4	1326.8	2384.40
4	1335.1	2477.50

Table 1: The observed response (weight gain) for a given treatment and value of the covariate (feed intake). Source: <http://www.stat.wisc.edu/~yandell/pda/>.

We use the following linear regression model for this data:

$$Y_{ij} = \mu_i + \tilde{\beta}_i(x_{ij} - \bar{x}_{..}) + \varepsilon_{ij}$$

where Y_{ij} is the response of the j th experimental unit ($j = 1, \dots, 3$) that is receiving the i th treatment ($i = 1, \dots, 4$), when the covariate takes the value x_{ij} . The ε_{ij} are independent and identically $N(0, \sigma^2)$ distributed and σ^2 is an unknown positive parameter. The μ_i and $\tilde{\beta}_i$ ($i = 1, \dots, 4$) are unknown parameters. Also, $\bar{x}_{..}$ denotes the mean of the x_{ij} ($i = 1, \dots, 4; j = 1, \dots, 3$).

We express this model in the form $Y = X\beta + \varepsilon$, where

$$Y = (Y_{11}, Y_{12}, Y_{13}, Y_{21}, Y_{22}, Y_{23}, Y_{31}, Y_{32}, Y_{33}, Y_{41}, Y_{42}, Y_{43})$$

$$\varepsilon = (\varepsilon_{11}, \varepsilon_{12}, \varepsilon_{13}, \varepsilon_{21}, \varepsilon_{22}, \varepsilon_{23}, \varepsilon_{31}, \varepsilon_{32}, \varepsilon_{33}, \varepsilon_{41}, \varepsilon_{42}, \varepsilon_{43}),$$

$\beta = (\mu_1, \mu_2, \mu_3, \mu_4, \tilde{\beta}_1, \tilde{\beta}_2, \tilde{\beta}_3, \tilde{\beta}_4)$ and X is the obvious 12×8 matrix.

As considered by Yandell (1997, p.271), we carry out a preliminary test of the null hypothesis $H_0 : \tilde{\beta}_1 = \tilde{\beta}_2 = \tilde{\beta}_3 = \tilde{\beta}_4$ against the alternative hypothesis H_A that the $\tilde{\beta}_i$ are not all the same, using an F test. Suppose that we use a 0.05 level of significance for this preliminary test. We express H_0 as $\tau = 0$ and H_A as $\tau \neq 0$, where $\tau = C^T \beta$ and

$$C^T = \begin{bmatrix} 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 & 1 \end{bmatrix}$$

Define the parameter of interest θ as follows. Let Y_1^* and Y_2^* denote the responses of two experimental units, receiving treatments 1 and 2 respectively, for the same value x^* of the covariate. In other words,

$$\begin{aligned} Y_1^* &= \mu_1 + \tilde{\beta}_1(x^* - \bar{x}_{..}) + \varepsilon_1^* \\ Y_2^* &= \mu_2 + \tilde{\beta}_2(x^* - \bar{x}_{..}) + \varepsilon_2^* \end{aligned}$$

where ε_1^* and ε_2^* are independent and identically $N(0, \sigma^2)$ distributed random variables. Let $\theta = E(Y_1^*) - E(Y_2^*) = \mu_1 - \mu_2 + (\tilde{\beta}_1 - \tilde{\beta}_2)(x^* - \bar{x}_{..})$. Thus, $\theta = a^T \beta$ where $a = (1, -1, 0, 0, (x^* - \bar{x}_{..}), -(x^* - \bar{x}_{..}), 0, 0)$. We suppose that $x^* - \bar{x}_{..} = 125.39$, which is the maximum value of $|x_{ij} - \bar{x}_{..}|$ for the data.

For this situation, $\|b\| = 0.96869$ and so the minimum coverage probability of the naive 0.95 confidence interval is 0.0846. This shows that this confidence interval is completely inadequate, in this situation.

5. Discussion

The poor coverage properties of naive confidence intervals found in this paper are presaged by the poor coverage properties of naive confidence intervals found in the context of a preliminary best subset variable selection by minimizing an AIC-type criterion, see e.g. Kabaila (2005), Kabaila & Leeb (2006) and Kabaila & Giri (2009). Apart from the form of preliminary model selection used, minimum AIC versus an F test, these papers differ from the present paper in that the present paper provides a method for computing the minimum coverage probability, whereas Kabaila (2005), Kabaila & Leeb (2006) and Kabaila & Giri (2009) provide only upper bounds on the minimum coverage probability of the naive confidence interval.

Appendix A: Proof of Theorem 1

In this appendix, we prove Theorem 1. Define $G = (\hat{\Theta} - \theta)/(\sigma\sqrt{v_{11}})$ and $H = (1/\sigma)V_{22}^{-1/2}\hat{\tau}$. Let f_H denote the probability density function of H . Note that

$$\begin{bmatrix} G \\ H \end{bmatrix} \sim N \left(\begin{bmatrix} 0 \\ \gamma \end{bmatrix}, \begin{bmatrix} 1 & b^T \\ b & I_s \end{bmatrix} \right). \quad (5)$$

where I_s denotes the $s \times s$ identity matrix. Thus the distribution of G , conditional on $H = h$, is $N(b^T(h - \gamma), 1 - \|b\|^2)$. Note that $(G, H^T)^T$ and W are independent random vectors. We use the notation

$$\chi(\mathcal{A}) = \begin{cases} 1 & \text{if } \mathcal{A} \text{ is true} \\ 0 & \text{if } \mathcal{A} \text{ is false} \end{cases}$$

where $\mathcal{A}$ is an arbitrary statement. This is similar to the Iverson bracket notation (Knuth, 1992).

By the law of total probability, the coverage probability of the naive $1 - \alpha$ confidence interval is

$$P(\theta \in I, F > \ell) + P(\theta \in J, F \leq \ell).$$

We divide the remainder of the proof into 2 parts.

Part 1: expression for $P(\theta \in I, F > \ell)$

Suppose that $\|\gamma\| > 0$. We prove the validity of the expressions (3) and (4) for $P(\theta \in I, F > \ell)$. The proofs of the validity of the other expressions for $P(\theta \in I, F > \ell)$ (given in Theorem 1) are similar and are omitted, for the sake of brevity.

Now $\{\theta \in I\} = \{-t(m)W \leq G \leq t(m)W\}$ and $F = H^T H / (sW^2)$. Thus

$$\begin{aligned} P(\theta \in I, F > \ell) &= P \left(-t(m)W \leq G \leq t(m)W, \frac{H^T H}{sW^2} > \ell \right) \\ &= \int \cdots \int \int_0^\infty P \left(-t(m)W \leq G \leq t(m)W, \frac{H^T H}{sW^2} > \ell \mid W = w, H = h \right) f_W(w) dw f_H(h) dh \\ &= \int \cdots \int \int_0^\infty P \left(-t(m)w \leq G \leq t(m)w \mid H = h \right) \chi(h^T h > s \ell w^2) f_W(w) dw f_H(h) dh \end{aligned} \quad (6)$$

Note that

$$\begin{aligned} &P(-t(m)w \leq G \leq t(m)w \mid H = h) \\ &= P(-t(m)w - b^T(h - \gamma) \leq Z \leq t(m)w - b^T(h - \gamma)) \quad \text{where } Z \sim N(0, 1 - \|b\|^2) \\ &= i(-b^T(h - \gamma), w; \|b\|). \end{aligned}$$

Thus

$$\begin{aligned}
(6) &= \int \cdots \int \int_0^\infty i(-b^T(h - \gamma), w; \|b\|) \chi(h^T h > s \ell w^2) f_W(w) dw f_H(h) dh \\
&= E\left(i(-b^T(H - \gamma), W; \|b\|) \chi(H^T H > s \ell W^2)\right) \\
&= E\left(i(-b^T(H - \gamma), W; \|b\|) \chi(W \leq \sqrt{H^T H / s \ell})\right). \tag{7}
\end{aligned}$$

We now find a simple formula for this expected value. Since $H \sim N(\gamma, I_s)$, $H = \gamma + RU$ where R is a nonnegative random variable and U is a random s -vector with the following distributions. The random vectors R^2 and U are independent, with $R^2 \sim \chi_s^2$ and U is distributed uniformly on the surface of the unit sphere in $\mathbb{R}^s$. Thus $b^T(H - \gamma) = R\|b\|L_b$, where $L_b = u_b^T U$. Also, $H^T H = \|\gamma\|^2 + 2\|\gamma\|RL_\gamma + R^2$, where $L_\gamma = u_\gamma^T U$. Hence

$$(7) = E\left(i(-\|b\|RL_b, W; \|b\|) \chi(W \leq \sqrt{(\|\gamma\|^2 + 2\|\gamma\|RL_\gamma + R^2)/s\ell})\right).$$

Note that (L_γ, L_b) , R and W are independent random vectors. Define the random vector (T_1, T_2) to be such that T_1 , T_2 , R and W are independent and T_1 and T_2 have the probability density functions f_{T_1} and f_{T_2} respectively, defined in Section 3. Define the unit s -vectors e_γ and e_b as follows. The vector e_γ has 1 as its first component and zeros for the remaining components. The vector e_b has first component ψ , second component $\sqrt{1 - \psi^2}$ and zeros for the remaining components. Because U is distributed uniformly on the surface of the unit sphere in $\mathbb{R}^s$, (L_γ, L_b) has the same distribution as $(e_\gamma^T U, e_b^T U)$. It follows from Fang and Wang (1994, p.49, pp.306–306 and p.308) that (U_1, U_2) has the same distribution as (a) $(\cos(2\pi T_1), \sin(2\pi T_1))$ for $s = 2$, (b) $(\cos(\pi T_1), \sin(\pi T_1) \cos(2\pi T_2))$ for $s = 3$ and $(\cos(\pi T_1), \sin(\pi T_1) \cos(\pi T_2))$ for $s > 3$. Thus, (L_γ, L_b) has the same distribution as

$$\left(\cos(2\pi T_1), \psi \cos(2\pi T_1) + \sqrt{1 - \psi^2} \sin(2\pi T_1)\right)$$

for $s = 2$,

$$\left(\cos(\pi T_1), \psi \cos(\pi T_1) + \sqrt{1 - \psi^2} \sin(\pi T_1) \cos(2\pi T_2)\right).$$

for $s = 3$ and

$$\left(\cos(\pi T_1), \psi \cos(\pi T_1) + \sqrt{1 - \psi^2} \sin(\pi T_1) \cos(\pi T_2)\right).$$

for $s > 3$. Thus $P(\theta \in I, F > \ell)$ is

$$E\left(i(-\|b\| R k(T_1; \psi), W; \|b\|) \chi(W \leq \sqrt{d(T_1, R; 2, \|\gamma\|)/s\ell})\right)$$

for $s = 2$ and

$$E\left(i(-||b|| Rk(T_1, T_2; s, \psi), W; ||b||) \chi(W \leq \sqrt{(d(T_1, R; s, ||\gamma||)/s\ell)})\right) \quad (8)$$

for $s \geq 3$. This leads to the expressions (3) and (4) for $P(\theta \in I, F > \ell)$ given in the theorem.

Part 2: expression for $P(\theta \in J, F \leq \ell)$

The derivation of the expression for $P(\theta \in J, F \leq \ell)$ is based on (1) and the fact that Θ^* , $\hat{\tau}$ and $R(\hat{\beta})$ are independent random vectors. Define $Q = (1/\sigma^2)\hat{\tau}^T V_{22}^{-1} \hat{\tau}$ and note that Q has a noncentral chi squared distribution with s degrees of freedom and noncentrality parameter $||\gamma||^2$. Note that

$$\{\theta \in J\} = \left\{ Z \in \left[b^T \gamma \pm t(m+s) \sqrt{\frac{mW^2 + Q}{m+s}} \sqrt{1 - ||b||^2} \right] \right\}$$

where Z , Q and W are independent random variables and $Z \sim N(0, 1 - ||b||^2)$. Also

$$F = \frac{\hat{\tau}^T V_{22}^{-1} \hat{\tau} / s}{\hat{\Sigma}^2} = \frac{Q/s}{W^2}.$$

Thus

$$\begin{aligned} P(\theta \in J, F \leq \ell) &= P\left(Z \in \left[b^T \gamma \pm t(m+s) \sqrt{\frac{mW^2 + Q}{m+s}} \sqrt{1 - ||b||^2} \right], Q \leq s\ell W^2\right) \\ &= \int_0^\infty \int_0^{s\ell w^2} j(||b|| ||\gamma|| \psi, q, w; ||b||) f_Q(q) f_W(w) dq dw \end{aligned}$$

by a method similar to that used in Part 1.

Appendix B: Numerical evaluation of the integrals in Theorem 1

We evaluate the integrals (2) and (3) in the statement of Theorem 1 as follows. We approximate (2) by

$$\int_0^{c_1} \int_0^{s\ell w^2} j(||b|| ||\gamma|| \psi, q, w; ||b||) f_Q(q) f_W(w) dq dw \quad (9)$$

for an appropriately chosen value of c_1 . We bound the error of this approximation as follows. Since $j(||b|| ||\gamma|| \psi, q, w; ||b||)$ is a probability,

$$\begin{aligned} 0 \leq (2) - (9) &\leq \int_{c_1}^\infty \int_0^{s\ell w^2} f_Q(q) f_W(w) dq dw \\ &\leq \int_{c_1}^\infty f_W(w) dw \\ &= P(M > m c_1^2) \quad \text{where } M \sim \chi_m^2. \end{aligned}$$

We choose c_1 sufficiently large that the right hand side is less than, say, 10^{-5} .

To evaluate (9), we transform the region of integration to a rectangle as follows. Change the variable of integration q in (9) to $q^* = q/(s\ell w^2)$, so that (9) is equal to

$$\int_0^{c_1} \int_0^1 j(\|b\| \|\gamma\| \psi, s\ell w^2 q^*, w; \|b\|) f_Q(s\ell w^2 q^*) s\ell w^2 f_W(w) dq^* dw.$$

The integrand is a smooth function of $(q^*, w) \in [0, 1] \times [0, c_1]$ and so this double integral is easily evaluated by numerical integration.

We approximate (3) by

$$\int_0^1 \int_0^{c_2} \int_0^u i(-\|b\| r k(t_1; \psi), w; \|b\|) f_W(w) f_R(r) dw dr dt_1 \quad (10)$$

for an appropriately chosen value of c_2 . We bound the error of this approximation as follows. Since $i(-\|b\| r k(t_1; \psi), w; \|b\|)$ is a probability,

$$\begin{aligned} 0 \leq (3) - (10) &\leq \int_0^1 \int_{c_2}^\infty \int_0^u f_W(w) dw f_R(r) dr f_T(t) dt \\ &\leq \int_{c_2}^\infty f_R(r) dr \\ &= P(R^2 > c_2^2) \quad \text{where } R^2 \sim \chi_s^2. \end{aligned}$$

We choose c_2 sufficiently large that the right hand side is less than, say, 10^{-5} .

To evaluate (10), we transform the region of integration to a rectangle as follows. Change the variable of integration w in (10) to $w^* = w/\sqrt{d(t_1, r; s, \|\gamma\|)/s\ell}$, so that (10) is equal to

$$\int_0^1 \int_0^{c_2} \int_0^1 i(-\|b\| r k(t_1; \psi), \sqrt{d(t_1, r; s, \|\gamma\|)/s\ell} w^*; \|b\|) f_W(\sqrt{d(t_1, r; s, \|\gamma\|)/s\ell} w^*) \sqrt{d(t_1, r; s, \|\gamma\|)/s\ell} f_R(r) dw^* dr dt_1.$$

The integrand is a smooth function of $(w^*, r, t_1) \in [0, 1] \times [0, c_2] \times [0, 1]$ and so this triple integral is easily evaluated by numerical integration.

The evaluation of the other integrals in the statement of Theorem 1 is similar to the evaluation of the integrals (2) and (3). The evaluation of (4) using MATLAB requires special comment. In MATLAB, the highest dimensional integral that one can evaluate using a built-in MATLAB function is a triple integral. We evaluate the quadruple integral (4) using MATLAB as follows. As before, let $u = \sqrt{d(t_1, r; s, \|\gamma\|)/s\ell}$. Define

$$g(w^*, t_1, r) = \int_0^1 i(-\|b\| r k(t_1, t_2; s, \psi), u w^*; \|b\|) f_{T_2}(t_2) dt_2$$

The integrand on the right-hand-side is a very smooth function of t_2 . We evaluate $g(w^*, t_1, r)$, to a good approximation, using a compound Simpson's rule with a specified number of subdivisions of the interval $[0, 1]$. The quadruple integral (4) is approximated by

$$\int_0^{c_2} \int_0^1 \int_0^1 g(w^*, t_1, r) f_W(u w^*) u f_{T_1}(t_1) f_R(r) dw^* dt_1 dr$$

which is evaluated using the MATLAB built-in function `triplequad`.

References

- Chin, S.F., Storkson, J.M., Albright, K.J., Cook, M.E. & Pariza, M.W.: Conjugate linoleic acid is a growth factor for rats as shown by enhanced weight gain and improved feed efficiency. *Journal of Nutrition* 124, 2344 – 2349 (1994)
- Fang, K.-T. & Wang, Y.: *Number-theoretic Methods in Statistics*. Chapman & Hall, London (1994)
- Farchione, D.: Interval estimators that utilize uncertain prior information. Unpublished Ph.D. thesis, Department of Mathematics and Statistics, La Trobe University (2009)
- Freund, R.J., Wilson, W.J. & Sa, P.: *Regression Analysis: Statistical Modeling of a Response Variable*, 2nd ed.. Elsevier, Academic Press, Burlington, Mass. (2006)
- Graybill, F.A.: *Theory and Application of the Linear Model*. Duxbury, Pacific Grove, CA (1976)
- Kabaila, P.: On the coverage probability of confidence intervals in regression after variable selection. *Australian & New Zealand Journal of Statistics* 47, 549–562 (2005).
- Kabaila, P., Leeb, H.: On the large-sample minimal coverage probability of confidence intervals after model selection. *Journal of the American Statistical Association* 101, 619–629 (2006)
- Kabaila, P., Giri, K.: Upper bounds on the minimum coverage probability of confidence intervals in regression after model selection. *Australian & New Zealand Journal of Statistics* 51, 271 – 288 (2009)
- Knuth, D.E.: Two notes on notation. *American Mathematical Monthly* 99, 403–422 (1992)
- Kuehl, R.O.: *Design of Experiments: Statistical Principles of Research Design and Analysis*, 2nd ed.. Brooks/Cole, Pacific Grove, CA (2002)

- Milliken, G.A., Johnson, D.E.: Analysis of Messy Data, Volume III: Analysis of Covariance. Chapman & Hall/CRC, Boca Raton, FL. (2002)
- Yandell, B.S.: Practical Data Analysis for Designed Experiments. Chapman & Hall, London, New York (1997).